

Codling moth (*Cydia pomonella*): Control methods in organic fruit production

Problem

Codling moth (*Cydia pomonella*) is the primary pest in organic fruit growing. It damages apple, pear, quince, rowan, medlar, walnut and kaki (photo A-D). Generations/years vary depending on the soil and climates zones.

Solution

Several control methods can be applied in organic farming depending on the infection rate, but plant protection products and mating disruption are the most common.

Benefits

Choosing suitable methods may reduce the time for intervention in the field. In the following season, precautionary measures can significantly reduce the spread of this pest.

Practical recommendation

- **Precautionary measures:** Remove woodpiles from the orchard and the affected fruit during thinning.
- **Mating disruption with pheromone dispensers (passive dispensers and aerosols):** Effective if used on a large scale and if the impact of the attack is limited. It is recommended to combine with other measures. Monitor the presence of the pest with traps set up in the orchard (Picture F) ([Practice abstract coming soon](#)).
- **Granulosis virus (CpGV):** Generally effective when the attack is mild and not sufficient to contain spreading in the case of stronger attacks. Treatments are best conducted during the twilight hours (Weblink 6)
- **Spinosine:** The most effective plant protection product available. It can be applied a maximum three times per year ([Practice abstract coming soon](#)).
- **Alt'Carpo nets:** Involves netting the plants with a mesh tighter than the average anti-hail nets. The netting can be done using a single-plot or single-row system. It is the most effective method to prevent adults from fleeing (Picture E).
- Entomopathogenic nematodes of the species *Steinernema feltiae* can parasitise codling moth larvae during the overwintering phase with an efficiency of up to 50%. Apply entomopathogenic nematodes with a high-water supply and high moisture content and a temperature (during treatment and the following 3 hours) of at least 10°C. On the day of treatment, the minimum temperature must not go below 0°C ([Practice abstract coming soon](#)).

Applicability box

Theme

Crop production, disease and pest control

Keywords

Temperate fruits, plant protection, pest control, biological control, plant protection product, preventive measures, mating disruption, physical barriers.

Context

Northern and central Europe.

Time of application and index of use

Index of use is the weighted average between time, cost and effectiveness expressed within a range of 0-5 (where 5 is the most effective and 0 the least effective measure).

-Mating disruption: before start of flight of first-generation; start of petal dropping.

(Index of use: 4.5)

-Granulosis-virus: during the first generation, when the first eggs begin to hatch.

(Index of use: 2,5)

-Spinosad: during first larvae detection.


(Index of use: 2.5)

-Nematodes: during autumn from September to October.

(Index of use: 2)

-Alt-carpo nets: after flowering.

(Index of use: 4)



Picture A-D. The caterpillar starts to penetrate the epidermis of the fruit (A) by digging tunnels into the pulp (B) until it reaches the central area where the seeds are located. As

a result of the trophic activity of the larvae, the fruits are filled with redness and excrement (C). At the end of its growth, the caterpillar emerges from the fruit by digging an exit tunnel and pupates on the woody organs (D).



Picture E: Alt'Carpo nets, a single-row system is shown. Picture F: To determine the percentage of damage before mating disruption treatments are applied, traps are placed into the orchard.

© A-D: Claudio Casera, E: Thomas Holtz, F: Josef Telfser. Organic farming team- Research Centre Laimburg (RCL).

Further information

Further reading

- M. Kelderer, C. Casera, E. Lardscheider, A. Rainer (2010). Controlling codling moth with different netting structures and their influence on crop yield and quality. 14th International Conference in Organic Fruit-Growing – Eco-fruit, Fördergemeinschaft Ökologischer Obstbau e. V. Weinsberg, Stuttgart, Deutschland, 183-190. Available here.
- E. Fritsch, K. Undorf-Spahn, J. Kienzle, J. Zimmer, B. Benduhn, C. Adolphi, C.P.W. Zebitz, J.A. Jehle (2020). Monitoring codling moth resistance to *Cydia pomonella* granulovirus (CpGV) in organic fruit growing in Germany. Proceedings ecofruit Conference 2020. Available here

Weblinks (Coming soon)

1. Practice abstract: Decision supporting systems to improve direct control methods of codling moth, IO
2. Practice abstract: Mating Disruption: The Key element of a successful building block strategy against *Cydia pomonella* in organic apple production, FOKO
3. Practice abstract: Innovative methods to reduce codling moth damage, IO
4. Practice abstract: Use of Nematodes in Autumn as preventive measure to reduce infestation level of *Cydia pomonella* for the following year, FOKO
5. Practice abstract: Avoiding bamboo sticks: Indirect measures to reduce infestation pressure of *Cydia pomonella* in organic pomme fruit production, FOKO
6. Practice abstract: Use of Caprovirusine-Products against *Cydia pomonella* in organic fruit growing to prevent resistance building of the pest, FOKO
7. Practice abstract: Codling moth: How to enhance beneficial insects in pome orchards, GRAB
8. Practice abstract: Codling moth on pear, DELPHY

About this practice abstract and

Publisher(s): Research Centre Laimburg- Italy

Laimburg 6, I-39040 Post Auer (BZ), Italy.

Tel. +39 0471 969500, <http://www.laimburg.it/>

Author: Alfredo Mora Vargas, Markus Kelderer

Contact: alfredo.moravargas@laimburg.it

Review: Ambra De Simone, Lauren Dietemann

Permalink: [Organic-farmknowledge.org/tool/xxx](https://organic-farmknowledge.org/tool/xxx)

Project name: BIOFRUITNET- Boosting Innovation in ORGANIC FRUIT production through stronger networks

Project website: <https://biofruitnet.eu>

© 2022

Codling moth (*Cydia pomonella*): Control methods in organic fruit production, Research Center Laimburg, Practice Abstract

This project has received funding from the European Union's Horizon 2020 research and innovation programme under (grant agreement No 862850). This communication only reflects the author's view. The Research Executive Agency is not responsible for any use that may be made of the information provided. The authors and editors do not assume responsibility or liability for any possible factual inaccuracies or damage resulting from the application of the recommendations in this practice abstract.

